

< 発表論文（下線は受賞者、責任者に「\*」） >

1. Kajiwara, C., Fumoto, K., Kimura, H., Nojima, S., Asano, K., Odagiri, K., Yamasaki, M., Hikita, H., Takehara, T., Doki, Y., Morii, E., and \*Kikuchi, A. p63-dependent Dickkopf3 expression promotes esophageal cancer cell proliferation via CKAP4. *Cancer Res.* doi: 10.1158/0008-5472.CAN-18-1749, 2018
2. Shinno, N., Kimura, H., Sada, R., Takiguchi, S., Mori, M., Fumoto, K., and Doki, Y., and \*Kikuchi, A. Activation of the Dickkopf1-CKAP4 pathway is associated with poor prognosis of esophageal cancer and anti-CKAP4 antibody may be a new therapeutic drug. *Oncogene* doi: 10.1038/s41388-018-0179-2, 2018
3. Fumoto, K., Takigawa-Imamura, H., Sumiyama, K., Kaneiwa, T., and \*Kikuchi, A. Modulation of apical constriction by Wnt signaling is required for lung epithelial shape transition. *Development* 144, 151-162, 2017
4. Kimura, H., Fumoto, K., Shojima, K., Nojima, S., Osugi, Y., Tomihara, H., Eguchi, H., Shintani, Y., Endo, E., Inoue, M., Doki, Y., Okumura, M., Morii, E., and \*Kikuchi, A. CKAP4 is a Dickkopf1 receptor and is involved in tumor progression. *J. Clin. Invest.* 126, 2689-2705, 2016
5. Matsumoto, S., Kurimoto, T., Taketo, M., Fujii, S., and \*Kikuchi, A. Wnt-Myb pathway suppresses KIT expression to control the timing of salivary proacinar differentiation and duct formation. *Development* 143, 2311-2324, 2016
6. Matsumoto, S., Fujii, S., Sato, A., Ibuka, S., Kagawa, Y., Ishii, M., and \*Kikuchi, A. A combination of Wnt and growth factor signaling induces Arl4c expression to form epithelial tubular structures. *EMBO J.* 33, 702-718, 2014
7. Naito, A., Sumida, T., Nomura, S., Liu, M.L., Higo, T., Nakagawa, A., Okada, K., Sakai, T., Hashimoto, A., Hara, Y., Shimizu, I., Zhu, W., Toko, H., Katada, A., Akazawa, H., Oka, T., Lee, J.K., Minamino, T., Nagai, T., Walsh, K., Kikuchi, A., Matsumoto, M., Botto, M., Shiojima, I., \*Komuro, I. Complement c1q activates canonical wnt signaling and promotes aging-related phenotypes. *Cell* 149, 1298-313, 2012
8. Kikuchi, K., Niikura, Y., Kitagawa, K., and \*Kikuchi, A. Dishevelled, a Wnt

- signaling component, is involved in mitotic progression with Plk1. *EMBO J.* 29, 3470-3483, 2010
9. Matsumoto, S., Fumoto, K., Okamoto, T., Kaibuchi, K., and \*Kikuchi, A. Binding of APC and disheveled mediates Wnt5a-regulated focal adhesion dynamics in migrating cells. *EMBO J.* 29, 1192-1204, 2010
10. Sato, A., Yamamoto, H., Sakane, H., Koyama, H., and \*Kikuchi, A. Wnt5a regulates distinct signaling pathways by binding to Frizzled2. *EMBO J.* 29, 41-54, 2010
11. Fumoto, K., Kadono, M., Izumi, N., and \*Kikuchi, A. Axin localizes to the centrosome and is involved in microtubule nucleation. *EMBO R.* 10, 606-613, 2009
12. Yamamoto, H., Kitadai, Y., Yamamoto, H., Oue, N., Ohdan, H., Yasui, W., and \*Kikuchi, A. Laminin  $\gamma$ 2 mediates Wnt5a-induced invasion of gastric cancer cells. *Gastroenterology* 137, 242-252, 2009
13. Yamamoto, H., Sakane, H., Yamamoto, H., Michiue, T., and \*Kikuchi, A. Wnt3a and Dkk1 regulate distinct internalization pathways of LRP6 to tune the activation of  $\beta$ -catenin signaling. *Dev. Cell* 15, 37-48, 2008
14. Zhu, W., Shiojima, I., Ito, Y., Li, Z., Ikeda, H., Yoshida, M., Naito, A., Nishi, J., Ueno, H., Umezawa, A., Minamino, T., Nagai, T., Kikuchi, A., Asashima, M., and \*Komuro, I. IGFBP-4 is a canonical Wnt inhibitor that promotes cardiac myogenesis. *Nature* 454, 345-349, 2008
15. Yamamoto, H., Komekado, H., and \*Kikuchi, A. Caveolin is necessary for Wnt-3a-induced internalization of LRP6 and accumulation of  $\beta$ -catenin. *Dev. Cell* 11, 213-223, 2006
16. Fumoto, K., Hoogenraad, C. C., and \*Kikuchi, A. GSK-3 $\beta$ -regulated interaction of BICD with dynein is involved in microtubule anchorage at centrosome. *EMBO J.* 25, 5670-5682, 2006
17. Kurayoshi, M., Oue, N., Yamamoto, H., Kishida, M., Inoue, A., Asahara, T., Yasui, W., and \*Kikuchi, A. Expression of Wnt-5a is correlated with aggressiveness of gastric cancer by stimulating cell migration and invasion. *Cancer Res.* 66, 10439-10448, 2006
18. Yamamoto, H., Ihara, M., Matsuura, Y., and \*Kikuchi, A. Sumoylation is

involved in  $\beta$ -catenin-dependent

activation of Tcf-4. *EMBO J.* 22, 2047-2059, 2003

19. Nakashima, S., Morinaka, K., Koyama, S., Ikeda, M., Kishida, M., Okawa, K., Iwamatsu, A., Kishida, S., and

\*Kikuchi, A. Small G protein Ral and its downstream molecules regulate endocytosis of EGF and insulin receptors.

*EMBO J.* 18, 3629-3642, 1999

20. Kitagawa, M., Hatakeyama, S., Shirane, M., Matsumoto, M., Ishida, N., Hattori, K., Nakamichi, I., Kikuchi, A.,

Nakayama, K-I., and \*Nakayama, K. An F-box protein, FWD1, mediates ubiquitin-dependent proteolysis of

$\beta$ -catenin. *EMBO J.* 18, 2401-2410, 1999

21. Ikeda, S., Kishida, S., Yamamoto, H., Murai, H., Koyama, S., and \*Kikuchi, A. Axin, a negative regulator of the

Wnt signaling pathway, forms a complex with GSK-3 $\beta$  and  $\beta$ -catenin and promotes GSK-3 $\beta$ -dependent

phosphorylation of  $\beta$ -catenin. *EMBO J.* 17, 1371-1384, 1998.